



<sup>1</sup> number between brackets: although for most cases no respiratory results were available at day 1, these are considered respiratory positive as this is inclusion criterion. However day 1 in the study can be multiple days later than the case was diagnosed with COVID-19 and therefore the respiratory result and feces result are not from approximately the same day.

Proportionally for persons with feces +ve at day 1 children have more frequently positive feces at day 15 compared to adults (Table 2).

**Table 2.** SARS-CoV-2 detection in feces of persons for which feces was collected at day 1 and day 15 by age group

Age group				
0 - <18 (n=117)			105 pairs	
			Day 1	
			+	-
	Day 15	+	16	4
		-	6	79
>=18 (n=123)			113 pairs	
			Day 1	
			+	-
	Day 15	+	15	0
		-	30	68

More than half of the children with +ve feces at day 1 showed no symptoms whereas at day 15 this is reversed, similar to in adults of whom most with +ve feces showed symptoms at day 1 and 15 (Table 3). For respiratory specimens at day 1 results are similar, of 25 positive children at day of specimen collection 14 did not show symptoms and for adults this was only 5 out of 27.

**Table 3.** SARS-CoV-2 detection in feces for persons showing symptoms or not; by age group

Age group	D1				D15			
0 - <18 (n=117)			109 pairs				106 pairs	
			Feces				Feces	
			+	-			+	-
	Symptoms	+	11	27	Symptoms	+	16	36
		-	13	58		-	5	49
>=18 (n=123)			118 pairs				111 pairs	
			Feces				Feces	
			+	-			+	-
	Symptoms	+	40	43	Symptoms	+	14	76
		-	7	28		-	1	20

As oral fluid has been suggested as a good alternative for respiratory specimens, especially for children the previously analysed oral fluid data [2], were combined with the current dataset and analysed (Table 4). In children who are positive in respiratory specimens feces outperforms oral fluid compared to adults. However, either way some of the respiratory positive persons would be missed using oral fluid and/or feces. However, as is clearly shown in table 1 also persons who test negative in respiratory specimens can test positive in feces. In the pilot study for oral fluid, none of the selected persons negative in respiratory specimens tested positive in oral fluid.[2].

**Table 4.** SARS-CoV-2 detection in feces and/or oral fluid of a selection of respiratory positive or negative persons at day 1 for which oral fluid was analysed; by age group

Age group	Respiratory positive				Respiratory negative			
0 - <18 (n=28)			21 pairs				6 pairs	
			Feces				Feces	
			+	-			+	-
	Oral fluid	+	12	1	Oral fluid	+	0	0
		-	5	3		-	0	6 <sup>1</sup>
>=18 (n=17)			15 pairs				2 pairs	
			Feces				Feces	
			+	-			+	-
	Oral fluid	+	12	2	Oral fluid	+	0	0
		-	0	1		-	1 <sup>2</sup>	1

<sup>1</sup> 2 are feces only positive at d14; 1 is respiratory and feces positive at d14

<sup>2</sup> is respiratory and feces positive at d14

In conclusion, feces as a convenient specimen for children, and especially young children, is a good alternative to taking nasopharyngeal and oropharyngeal swabs (or nasal aspirate). Although numbers are low, in comparison with collection of oral fluid feces seem to outperform oral fluid in positivity rate among children with positive respiratory specimens. We can also conclude that feces is a good addition to the diagnostic repertoire, especially if respiratory specimens test negative and the suspicion for COVID-19 is strong.

Proposal for specimens collection in young children at 'basisschool' age and below, depended on what a laboratory can process:

1. Collect feces
2. If possible collect also oral fluid. For the time being using Oracol sponge S10 or S14; several other systems, including those with nucleic acid preservative are investigated [3,4]. However, for very young children the sponge seems most convenient to use.

If testing capacity does not allow two specimens per child and if testing is used for contact investigation, feces is the preferred specimen.

