To: [redacted]@rivm.nl]; [redacted]@rivm.nl]; [redacted]@rivm.nl]; [redacted]@rivm.nl]; [redacted]@rivm.nl]; [redacted]@rivm.nl]; [redacted]@rivm.nl]; [redacted]@rivm.nl]
From: [redacted]
Sent: Fri 5/29/2020 8:22:54 AM
Subject: RE: % mensen dat SARS-CoV-2 in feces uitscheidt ?
Received: Fri 5/29/2020 8:22:55 AM
Wai-Shun Chan et al - 2020 - A systematic review on COVID-19 Urological manifestations, viral RNA.pdf

From: [redacted]
Sent: vrijdag 29 mei 2020 10:22
To: [redacted]@rivm.nl>; [redacted]@rivm.nl>; [redacted]@rivm.nl>; [redacted]@rivm.nl>; [redacted]@rivm.nl>; [redacted]@rivm.nl>; [redacted]@rivm.nl>; [redacted]@rivm.nl>
Subject: RE: % mensen dat SARS-CoV-2 in feces uitscheidt ?

Goed nieuws x2, heb even meeste recente literatuur gezocht en review gevonden. Gemiddelde over 21 studies:
65.82% (95% CI 45.71–83.51%) of COVID-19 patients had
positive viral RNA in stool samples, which were detected from 2 to 47 days from symptom onset.
Dus mooi voor de surveillance dat in ieder geval onder patienten % boven de 50% ligt.
En dat deel van antwoord er nu is. Vraag staat dan nog open hoe dit onder asymptomaten is.

Ik stuur straks nog aantekeningen van gisteren plus alle andere beloofde studies.

Groeten,

[redacted]

De volgende pagina is verwijderd i.v.m. blanco.