

Symptoms included in primary and secondary case definitions

Janssen ("moderate to severe/critical")

>=1 of:

Respiratory >=20
 Abnormal pulse ox but >93%
 Clinical/Radiological pneumonia
 DVT
 Shortness of breath/difficulty breathing

OR

>=2 of:

Fever >=38C
 Heart rate >=90
 Shaking chills
 Sore throat
 Cough
 Malaise
 Headache
 Myalgia
 GI sx (N/V/D/abdominal pain)
 Olfactory/taste disorder
 Red/bruised toes

Secondary

Mild COVID-19

At least 1 of sx above (except heart rate) or:

Chest congestion
 Runny nose
 Wheezing
 Skin rash
 Eye irritation/discharge

"FDA harmonized"

fever or chills
 cough
 shortness of breath/difficulty breathing
 fatigue
 muscle or body aches
 headache
 loss of taste or smell
 sore throat
 congestion

runny nose
nausea or vomiting
diarrhea

(note that the Janssen "FDA harmonized"
actually seems to align with other vaccines'
"expanded CDC symptom list"

| Moderna | Pfizer |
|--|--|
| <u>>=1 of:</u> | <u>>=1 of:</u> |
| Clinical/Radiological pneumonia | |
| Shortness of breath | Shortness of breath |
| Cough | |
| OR | |
| <u>>=2 of:</u> | |
| Fever >=38C | Fever >=38C |
| Chills | Chills |
| Sore throat | Sore throat |
| | Cough |
| Headache | |
| Myalgia | Muscle pain |
| | Diarrhea or vomiting |
| Olfactory/taste disorder | Loss of taste or smell |
| <u>At least 1 of above or expanded CDC symptom</u> | <u>At least 1 of above or expanded CDC symptom list:</u> |
| Fatigue | Fatigue |
| Nasal congestion | Nasal congestion/runny nose |
| Nausea/vomiting | |
| Diarrhea | |
| | Headache |
| | Nausea |

De volgende pagina is verwijderd ivm blanco

Symptoms included in primary and secondary case definitions

| | |
|------------------|--|
| | Janssen |
| Primary | Moderate/severe COVID-19 |
| | <u>>=1 of:</u> |
| | Respiratory >=20 |
| | Abnormal pulse ox but >93% |
| | Clinical/Radiological pneumonia |
| | DVT |
| | Shortness of breath/difficulty breathing |
| | OR |
| | <u>>=2 of:</u> |
| | Fever >=38C |
| | Heart rate >=90 |
| | Shaking chills |
| | Sore throat |
| | Cough |
| | Malaise |
| | Headache |
| | Myalgia |
| | GI sx (N/V/D/abdominal pain) |
| | Olfactory/taste disorder |
| | Red/bruised toes |
| Secondary | Mild COVID-19 |
| | <u>At least 1 of bolded sx above or:</u> |
| | Chest congestion |
| | Runny nose |
| | Wheezing |
| | Skin rash |
| | Eye irritation/discharge |
| "FDA harmonized" | fever or chills |
| | cough |
| | shortness of breath/difficulty breathing |
| | fatigue |
| | muscle or body aches |
| | headache |
| | loss of taste or smell |
| | sore throat |
| | congestion |
| | runny nose |
| | nausea or vomiting |

diarrhea

(note that the Janssen "FDA harmonized"
actually seems to align with other vaccines!
"expanded CDC symptom list"

| Moderna | Pfizer |
|--|--|
| Primary | Primary |
| <u>>=1 of:</u> | <u>>=1 of:</u> |
| Clinical/Radiological pneumonia | |
| Shortness of breath | Shortness of breath |
| Cough | |
| OR | |
| <u>>=2 of:</u> | |
| Fever >=38C | Fever >=38C |
| Chills | Chills |
| Sore throat | Sore throat |
| | Cough |
| Headache | |
| Myalgia | Muscle pain |
| | Diarrhea or vomiting |
| Olfactory/taste disorder | Loss of taste or smell |
| <u>At least 1 of above or expanded CDC symptom list:</u> | <u>At least 1 of above or expanded CDC symptom list:</u> |
| Fatigue | Fatigue |
| Nasal congestion | Nasal congestion/runny nose |
| Nausea/vomiting | |
| Diarrhea | |
| | Headache |
| | Nausea |

De volgende pagina is gelakt ivm blanco

| Janssen | Moderna | Pfizer |
|---|---|---|
| "Severe/Critical" | "Severe COVID-19" | "Severe COVID-19" |
| 1 of the following: | any of the following: | at least 1 of the following: |
| Resp rate ≥ 30 bpm | Resp rate ≥ 30 bpm | Resp rate ≥ 30 bpm |
| Heart rate ≥ 125 bpm | Heart rate ≥ 125 bpm | Heart rate ≥ 125 bpm |
| SpO2 93% | SpO2 93% | SpO2 93% |
| PaO2/FiO2s < 300 mmHg | PaO2/FiO2s < 300 mmHg | PaO2/FiO2s < 300 mmHg |
| Respiratory failure (needing high-flow O2, non-invasive ventilation, mechanical ventilation, or ECMO) | Respiratory failure or ARDS (needing high-flow O2, non-invasive ventilation, mechanical ventilation, or ECMO) | Respiratory failure (needing high-flow O2, non-invasive ventilation, mechanical ventilation, or ECMO) |
| Shock (SBP < 90 , DBP < 60 , or requiring vasopressors) | Shock (SBP < 90 , DBP < 60 , or requiring vasopressors) | Shock (SBP < 90 , DBP < 60 , or requiring vasopressors) |
| Significant acute renal, hepatic, or neurologic dysfunction | Significant acute renal, hepatic, or neurologic dysfunction | Significant acute renal, hepatic, or neurologic dysfunction |
| Admission to ICU | Admission to ICU | Admission to ICU |
| Death | Death | Death |

Janssen

| | |
|----------|---|
| | Redness |
| | Grade 3: >10.0 cm |
| Local | Grade 4: necrosis |
| | Swelling |
| | Grade 3: >10.0 cm |
| | Grade 4: necrosis or exfoliative dermatitis |
| | Pain at the injection site |
| | Grade 3: any use of narcotic pain reliever or prevents daily activity |
| | Grade 4: emergency room visit or hospitalization |
| | Fever |
| | Grade 3: >38.9 to 40 |
| Systemic | Grade 4: >40 |
| | Fatigue |
| | Grade 3: any use of narcotic pain reliever or prevents daily activity |
| | Grade 4: hospitalization or inability to perform basic self care |
| | Headache |
| | Grade 3: any use of narcotic pain reliever or prevents daily activity |
| | Grade 4: hospitalization or inability to perform basic self care |
| | Nausea |
| | Grade 3: prevents daily activity |
| | Grade 4: hospitalization or inability to perform basic self care |
| | Myalgia |
| | Grade 3: any use of narcotic pain reliever or prevents daily activity |
| | Grade 4: hospitalization or inability to perform basic self care |

Moderna

Redness

Grade 3: >10.0 cm

Grade 4: necrosis

Swelling

Grade 3: >10.0 cm

Grade 4: necrosis or exfoliative dermatitis

Pain at the injection site

Grade 3: any use of prescription pain reliever or prevents daily activity

Grade 4: emergency room visit or hospitalization

Axillary Swelling/Tenderness

Grade 3: any use of prescription pain reliever or prevents daily activity

Grade 4: emergency room visit or hospitalization.

Fever

Grade 3: >38.9 to 40

Grade 4: >40

Fatigue

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

Headache

Grade 3: any use of prescription pain reliever or prevents daily activity

Grade 4: emergency room visit or hospitalization

Chills

Grade 3: prevents daily activity, required medical intervention

Grade 4: emergency room visit or hospitalization

Nausea/Vomiting

Grade 3: prevents daily activity, requires intravenous hydration

Grade 4: emergency room visit or hospitalization

Myalgia

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

Arthralgia

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

Pfizer

Redness

Grade 3: >10.0 cm

Grade 4: necrosis

Swelling

Grade 3: >10.0 cm

Grade 4: necrosis or exfoliative dermatitis

Pain at the injection site

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

Fever

Grade 3: >38.9 to 40

Grade 4: >40

Fatigue

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

Headache

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

Chills

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

Vomiting

Grade 3: requires intravenous hydration

Grade 4: emergency room visit or hospitalization

Diarrhea

Grade 3: 6 or more loose stools in 24 hours

Grade 4: emergency room visit or hospitalization

New or worsening muscle pain

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization

New or worsening joint pain

Grade 3: prevents daily activity

Grade 4: emergency room visit or hospitalization