

## Aerogenic transmission of SARS-CoV-2

- Multiple parties, including the Ministry of Health, Welfare and Sport, asked the RIVM for advice about aerogenic transmission of SARS-CoV-2 over a longer distance than 1,5 meters and possible implications for ventilation inside buildings.
- Aerosols are formed during speech, coughing and sneezing. However, at this moment there is not enough evidence that aerogenic transmission plays a relevant role in the spreading of SARS-CoV-2.
- Based on the current insights, alterations to ventilation systems are not necessary. Current guidelines for ventilation can be used.
- There will be a separate advice on the role of aerosols in singing and sports.

---

## Aerogenic transmission of SARS-CoV-2

### *Research questions & methods:*

Can SARS-CoV-2 be transmitted through aerosols? If so, are there any additional measures necessary for ventilation systems? To answer these questions, relevant literature was searched through databases like PubMed and Embase. Only studies regarding SARS-CoV-2 were included. Literature specifically on singing, exercise and fecal-oral transmission was not included.

### *Results*

Aerosols are formed during speech, coughing and sneezing (Asadi et al., 2020). Under laboratory circumstances it has been shown that SARS-CoV-2 can survive in aerosols for several hours (Fears et al., 2020; Van Doremalen et al., 2020). However, an  $R_0$  of 2-4 does not indicate aerogenic transmission. Different studies within hospitals have found virus RNA in air samples (Liu et al., 2020; Ong et al., 2020; Santarpia et al., 2020). In contrast, two other studies did not find RNA in air samples, even when taken close to a patient's chin (Cheng et al., 2020; Faridi et al., 2020). In some occasions outside the hospital aerogenic transmission is named as a possible transmission route (Brurberg, 2020; Lu et al., 2020; Wang et al., 2020). However, in these studies other transmission routes cannot be excluded. There are no studies that have measured the presence of virus in the air in public spaces, such as schools and supermarkets. Research on the outbreak aboard the Diamond Princess cruise ship showed that transmission took place via direct contact and contaminated surfaces, and not through the air-conditioning system (Xu et al., 2020).

### *Conclusion*

Different studies based on epidemiologic, virologic and modelling research do not show aerogenic transmission as a main route for the spread of SARS-CoV-2. At this moment, there is not enough evidence that aerogenic transmission plays a relevant role in the spread of SARS-CoV-2 over longer distances. Based on the current insights, alterations to ventilation systems are not necessary. Current guidelines for ventilation can be used. The background document about this subject is published on the [RIVM website](#).

5.1.2e ( 5.1.2e @rivm.nl)  
5.1.2e ( 5.1.2e @rivm.nl)

### *References*

- Asadi, S., Bouvier, N., Wexler, A. S., & Ristenpart, W. D. (2020). The coronavirus pandemic and aerosols: Does COVID-19 transmit via expiratory particles?.
- Brurberg, K. G. SARS-CoV-2, MERS-CoV and SARS-CoV and risk of airborne transmission. Center for Disease Control and Prevention. *Person-to-person spread: 13 April 2020*. Center for Disease Control and Prevention.
- Cheng VCC, Wong SC, Chen JHK, et al. Escalating infection control response to the rapidly evolving epidemiology of the Coronavirus disease 2019 (COVID-19) due to SARS-CoV-2 in Hong Kong. *Infect Control Hosp Epidemiol*. 2020 Mar 5:1-24.
- ECDC. Coronavirus disease 2019 (COVID-19) in the EU/EEA and the UK– ninth update. 23 April 2020. [Link](#).
- Faridi, S., Niazi, S., Sadeghi, K., Naddafi, K., Yavarian, J., Shamsipour, M., ... & Momeniha, F. (2020). A field indoor air measurement of SARS-CoV-2 in the patient rooms of the largest hospital in Iran. *Science of The Total Environment*, 138401.
- Fears, A. C., Klimstra, W. B., Duprex, P., Hartman, A., Weaver, S. C., Plante, K. S., ... & Nalca, A. (2020). Comparative dynamic aerosol efficiencies of three emergent coronaviruses and the unusual persistence of SARS-CoV-2 in aerosol suspensions. *medRxiv*.
- Federatie Medisch Specialisten, Leidraad medische procedures die een infectieuze aerosol genereren (IAGP) met SARS-CoV-2, versie 1, 29 april 2020, [link](#).
- Kim, Y. I., Kim, S. G., Kim, S. M., Kim, E. H., Park, S. J., Yu, K. M., ... & Um, J. (2020). Infection and rapid transmission of sars-cov-2 in ferrets. *Cell host & microbe*.
- Landelijke Coördinatie Infectieziektebestrijding. *Richtlijn COVID-19: 21 maart 2020*. Rijksinstituut voor Volksgezondheid en Milieu.
- Lee, J., Yoo, D., Ryu, S., Ham, S., Lee, K., Yeo, M., ... & Yoon, C. (2019). Quantity, Size Distribution, and Characteristics of Cough-generated Aerosol Produced by Patients with an Upper Respiratory Tract Infection. *Aerosol and Air Quality Research*, 19(4), 840-853.
- [5.1.2e](#), Ning, Z., Chen, Y., Guo, M., [5.1.2e](#), Gali, N. K., ... & Liu, X. (2020). Aerodynamic characteristics and RNA concentration of SARS-CoV-2 aerosol in Wuhan hospitals during COVID-19 outbreak. *bioRxiv*.
- Lu, J., Gu, J., Li, K., Xu, C., Su, W., Lai, Z., ... & Yang, Z. (2020). COVID-19 Outbreak Associated with Air Conditioning in Restaurant, Guangzhou, China, 2020. *Emerging Infectious Diseases*, 26(7).
- Ong, S. W. X., Tan, Y. K., Chia, P. Y., Lee, T. H., Ng, O. T., Wong, M. S. Y., & Marimuthu, K. (2020). Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) from a symptomatic patient. *Jama*.
- Richard, M., Kok, A., de Meulder, D., Bestebroer, T.M., Lamers, M.M., Okba, N.M.A., Fentener van Vlissingen, M., Rockx, [5.1.2e](#), B.L., Koopmans, M.P.G., Fouchier, R.A.M., Herfst, S., 2020. SARS-CoV-2 is transmitted via contact and via the air between ferrets. *bioRxiv*, 2020.2004.2016.044503.
- Santarpià, J. L., Rivera, D. N., Herrera, V., Morwitzer, M. J., Creager, H., Santarpià, G. W., ... & Lawler, J. V. (2020). Transmission Potential of SARS-CoV-2 in Viral Shedding Observed at the University of Nebraska Medical Center. *medRxiv*.
- Tian Y, Rong L, Nian W, He Y (2020). Review article: gastrointestinal features in COVID-19 and the possibility of faecal transmission. *Aliment Pharmacol Ther*. 51(9):843-851.
- van Doremalen N., Bushmaker, T., Morris, D. H., Holbrook, M. G., Gamble, A., Williamson, B. N., ... & Lloyd-Smith, J. O. (2020). Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *New England Journal of Medicine*.

- Wang, J., & Du, G. (2020). COVID-19 may transmit through aerosol. *Irish Journal of Medical Science (1971-)*, 1-2.
- World Health Organization. (2020). *Modes of transmission of virus causing COVID-19: implications for IPC precaution recommendations: scientific brief, 29 March 2020* (No. WHO/2019-nCoV/Sci\_Brief/Transmission\_modes/2020.2). World Health Organization.
- Wölfel, R., Corman, V.M., Guggemos, W. *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* (2020). <https://doi.org/10.1038/s41586-020-2196-x>
- WHO technical brief (2020). Water, sanitation, hygiene and waste management for COVID-19. [WHO](#).
- World Health Organization. (2020). Infection prevention and control during health care when novel coronavirus (nCoV) infection is suspected Interim guidance, 19 March 2020. WHO.
- Xiao F, Tang M, Zheng X, [5.1.2e](#), [5.1.2e](#), Shan H (2020). Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology*. Mar 3. pii: S0016-5085(20)30282-1.
- Xu, P., Qian, H., Miao, T., Yen, H. L., Tan, H., Cowling, B. J., & Li, Y. J. (2020). Transmission routes of Covid-19 virus in the Diamond Princess Cruise ship. *medRxiv*.