


PIENTER-Corona:

Prospective serosurveillance study of SARS-CoV-2 in the general population of the Netherlands

Correcting seroprevalence estimates



(10)(2e) – Epidemiologist at National Institute for Public Health and the Environment (RIVM)



Design PIENTER-3 and PICO1

- **PIENTER-3:**
 - Nationwide sample of the Dutch population (2016/2017) to look into protection against VPD
 - Two-stage cluster design: six regions, comprising 49 randomly assigned municipalities
 - Biobank (including pre-sera) of 7600 participants
 - N=6102 participants (80%) gave consent to be approached in the future
- **PIENTER-Corona (PICO1):**
 - N=3207 (aged 2-90y, across the NL) provided a self-collected fingerstick blood sample and filled out a questionnaire on risk factors






Antibody testing and correcting

- **Step 1:** All PICO1-serum samples were tested for the presence of SARS-CoV-2 IgG antibodies - targeted at the S1-part of the spike protein - using our Multiplex immunoassay (Luminex technology)
 - **Step 2:** Due to the expected low seroprevalence in this epidemic phase a specificity-optimized cutoff value (99%) for seropositivity was determined, using a validation panel consisting of:
 - 115 PCR-positive samples (including mild and severe COVID-19 patients)
 - 400 controls (i.e., pre-pandemic samples, including a batch of ILI-samples (also HCoVs)), as well as from PIENTER-3 and PIENTER-2)
- > Sensitivity at this cutoff (2.37 AU/mL) was 84.4%




Antibody testing and correcting

- **Step 3:** Seropositive PICO1-samples and those 25% below the cutoff were retested (n=138) -> GMC was used for further statistical analyses
- **Step 4:** 129/138 PICO1-samples had a pre-pandemic PIENTER-3-sample, and these were tested to correct for false-positivity:
 - > We classified PICO1-samples with a seropositive pre-pandemic serum (based on our validated cutoff) as seronegative (blue lines) (n=26)





Antibody testing and correcting



Final overview: IgG antibody concentration (AU/mL (log)) against SARS-CoV-2 of all individual PICO1-samples, by age (years) (**left side**) and distributed by means of a histogram (**right side**).

-> Seropositive samples: **76**



Antibody testing and correcting

- **Step 5: Statistical correction:**

- **A:** Calculate apparent weighted prevalence (with e.g., 95% Wilson/ClopperPearson CIs):
 - Correct for the survey design: strata (i.e., regions) and clusters (i.e., municipalities)
 - Include weights to match the distribution of the general public in 2020; here based on sex, age, ethnic background and degree of urbanization
- **B:** Calculate true weighted prevalence:
 - Correct for test specifics via Rogan-Gladen estimator, with sensitivity of 84.4% and assuming a specificity of 100% after cross-validation with pre-sera, using formula:

$$\frac{awp + SP - 1}{SE + SP - 1} \longrightarrow \frac{0.023 + 1.0 - 1}{0.844 + 1.0 - 1} \longrightarrow \mathbf{2.8\% (2.1-3.7)}$$



Additionally: Smooth age-specific seroprevalence

-> Logistic regression in a Generalized Additive Model using penalized splines (mgcv package in R), with additional Rogan&Gladen correction

